

COMPLETE SUBGRAPHS OF THE GRAPHS OF CONVEX POLYTOPES

S. GALLIVAN, E.R. LOCKEBERG and P. MCMULLEN

Department of Mathematics, University College London, Gower Street, London WC1E 6BT, England

Received 31 July 1979

It is shown that if three vertices of the graph $G(P)$ of a convex 3-polytope P are chosen, then $G(P)$ contains a refinement of the complete graph C_4 on four vertices, for which the three chosen vertices are principal (that is, correspond to vertices of C_4 in the refinement). In general, all four vertices may not be preassigned as principal. For dimensions $d \geq 4$, simple (simplicial) d -polytopes are constructed whose graphs contain sets of three (four) vertices, which cannot all be principal in any refinement of C_{d+1} .

1. Introduction

The graph of a d -polytope (d -dimensional convex polytope) P is the complex $G(P)$ consisting of the vertices and edges of P . The complete graph C_n on n vertices is the graph with n vertices, every pair of which is joined by an edge. The graph of the d -simplex T^d is isomorphic to C_{d+1} . All the graphs we consider here may be regarded as embedded in some euclidean space.

A cell-complex B is a *refinement* of another complex A if there is a homeomorphism $f: B \rightarrow A$ such that, for each cell C of A , $f^{-1}(C)$ is a union of cells of B . The homeomorphism f is called a *refinement map* from B to A . A cell D of B is called *principal* if $f(D)$ is a cell of A . In the particular case of graphs, note that the vertices of a principal edge will themselves be principal. If G contains a subgraph which is a refinement of a graph H , we say that G contains a refinement of H .

Grünbaum ([1]; see also [2, p. 200]) has shown, as a corollary of a stronger result, that the graph $G(P)$ of a d -polytope P contains a refinement of C_{d+1} . Furthermore, if any edge E of P is chosen, a refinement of C_{d+1} may be found for which E , and hence also its vertices, are principal. Since C_{d+1} is isomorphic to $G(T^d)$, C_{d+1} is the largest complete graph which is a refinement in the graph of every d -polytope.

In this paper, we consider the problem of determining how many vertices of a d -polytope P can be preassigned as principal in such a refinement of C_{d+1} . Grünbaum's result implies that we can preassign as principal any vertex of P , or any two vertices if they happen to be endpoints of the same edge. Larman and

Mani [3] show that if P is simplicial, two vertices of P can be preassigned as principal, however they are situated (we shall discuss this in Section 3).

If P is a d -polytope, define $r(P)$ to be the largest positive integer k such that, if any k vertices of P are chosen, there is a refinement of C_{d+1} in $G(P)$ such that all the chosen vertices are principal. Further define

$$r(d) = \min\{r(P) \mid P \in \mathcal{P}^d\}$$

and

$$r_s(d) = \min\{r_s(P) \mid P \in \mathcal{P}_s^d\},$$

where \mathcal{P}^d (\mathcal{P}_s^d) is the family of all (simplicial) d -polytopes. Of course, $r(d) \leq r_s(d)$.

A 1-polytope is a line segment; hence $r(1) = r_s(1) = 2$.

The graph of a 2-polytope (convex polygon) is a simple circuit with n edges, for some $n \geq 3$. The complete graph C_3 is a circuit of three edges. It is thus clear that $G(P)$ is a refinement of C_3 , and three vertices can always be preassigned. Thus $r(2) = r_s(2) = 3$.

In Section 2 of this paper, we show that $r(3) = r_s(3) = 3$. In Section 3, we construct examples to show that for $d \geq 4$, $r(d) \leq 2$ and $r_s(d) \leq 3$. (Larman and Mani give in [3] an example of a simplicial 3-polytope P with $r(P) < 4$, but their supposed examples in higher dimensions are not simplicial.) The table below summarizes our present knowledge of $r(d)$ and $r_s(d)$. The authors' guess is that for $d \geq 4$, $r(d) = 2$ and $r_s(d) = 3$.

d	$r(d)$	$r_s(d)$
1	2	2
2	3	3
3	3	3
≥ 4	1 or 2	2 or 3

2. Three dimensions

Easy examples show that $r(3) \leq 3$; for instance, the four vertices of a square face of a triangular prism cannot all be principal. The main result of this paper shows that there is, in fact, equality here.

Theorem 1. *Let x_1, x_2 and x_3 be distinct vertices of a 3-polytope P . Then $G(P)$ contains a refinement of C_4 , for which x_1, x_2 and x_3 are three of the principal vertices.*

Proof. We prove Theorem 1 in two stages. We first show that there are three arcs in $G(P)$ from x_1 to x_2 , and disjoint except at x_1 and x_2 , one of which passes through x_3 . Since $G(P)$ is planar, we can think of $G(P)$ as embedded in the euclidean plane E^2 . Since $G(P)$ is 3-connected, there are arcs Γ_1, Γ_2 and Γ_3 in $G(P)$ joining x_1 and x_2 , and disjoint except at their endpoints. If $x_3 \in$

$\Gamma_1 \cup \Gamma_2 \cup \Gamma_3 = \Gamma$, say, we are done; thus we may assume that x_3 lies in one of the connected components of $E^2 \setminus \Gamma$, say the component bounded by $\Gamma_2 \cup \Gamma_3$. Since $G(P)$ is 3-connected, there exist arcs Λ_1, Λ_2 and Λ_3 in $G(P)$ from x_3 to $\Gamma_2 \cup \Gamma_3$, disjoint except at x_3 . We may suppose that for $i = 1, 2, 3$, $\Lambda_i \cap (\Gamma_2 \cup \Gamma_3)$ is a single point y_i . One of the arcs Γ_2 or Γ_3 must meet two of the arcs Λ_1, Λ_2 or Λ_3 ; without loss of generality, suppose that Γ_3 meets Λ_1 and Λ_2 , and that y_1 is nearer to x_1 on Γ_3 than y_2 . Writing $\Delta(u, v)$ for that part of the arc Δ between its two vertices u and v , we then see that we can replace Γ_3 by

$$\Gamma'_3 = \Gamma_3(x_1, y_1) \cup \Lambda_1 \cup \Lambda_2 \cup \Gamma_3(y_2, x_2),$$

noting that Γ'_3 meets Γ_1 or Γ_2 only in x_1 and x_2 .

Notice that this conclusion does not hold for general 3-connected graphs, such as the complete bipartite graph on two groups of three vertices.

So, changing Γ'_3 back to Γ_3 , we have arcs Γ_i ($i = 1, 2, 3$) from x_1 to x_2 in $G(P)$, disjoint except at their endpoints, with $x_3 \in \Gamma_3$. Using the 3-connectedness of $G(P)$ again, there are arcs Δ_i ($i = 1, 2, 3$) in $G(P)$, disjoint except at x_3 , leading from x_3 to $\Gamma_1 \cup \Gamma_2$. We may suppose the Δ_i so chosen, that the number of edges in $\Delta = \Delta_1 \cup \Delta_2 \cup \Delta_3$, which are not already in $\Gamma = \Gamma_1 \cup \Gamma_2 \cup \Gamma_3$, is minimal; that is, we introduce as few new edges as possible. Then we claim that each of x_1 and x_2 lies in Δ . For, if (say) $x_1 \notin \Delta$, let w_1 be the nearest vertex of Γ_3 to x_1 on Δ ; suppose that $w_1 \in \Delta_1$, and if $w_1 = x_3$, further suppose that $\Delta_1 \neq \Gamma_3(x_3, x_2)$. We can then replace Δ_1 by $\Delta_1(x_3, w_1) \cup \Gamma_3(w_1, x_1)$, which gives a contradiction to our minimality assumption. Thus $x_i \in \Delta$ ($i = 1, 2$), and it is now clear that $\Gamma_1 \cup \Gamma_2 \cup \Delta$ is the required refinement, which completes the proof of Theorem 1.

We have thus shown that $r(3) = 3$. The example of Theorem 4 in Section 3 will show that $r_s(3) \leq 3$, and so $r_s(3) = 3$ also.

3. Four or more dimensions

In contrast to Theorem 1, in dimensions higher than three we have the following result.

Theorem 2. *Let $d \geq 4$. Then there is a simple d -polytope P , with some three vertices which cannot all be principal for any refinement of C_{d+1} in $G(P)$.*

Proof. Choose any $p, q \geq 2$, with $p + q = d$, and let $P = T^p \times T^q$ be the cartesian product of a p -simplex and a q -simplex. If we write $\text{vert } T^p = \{x_0, \dots, x_p\}$ and $\text{vert } T^q = \{y_0, \dots, y_q\}$, then the vertices of P are all (x_i, y_j) , with $i = 0, \dots, p$ and $j = 0, \dots, q$. The two vertices (x_i, y_j) and $(x_{i'}, y_{j'})$ determine an edge of P if and only if $i = i'$ or $j = j'$, but not both.

Let us choose $z_0 = (x_0, y_0)$ and $z_1 = (x_1, y_1)$ as principal vertices. We shall show

that there are exactly two refinements of C_{d+1} in $G(P)$ with z_0 and z_1 principal, and that neither contains $z_2 = (x_2, y_2)$ as a principal vertex. Hence, for no refinement of C_{d+1} are z_0, z_1 and z_2 all principal.

Let the desired refinement of C_{d+1} with z_0 and z_1 principal be the subgraph B of $G(P)$. Since both T^p and T^q are simple, so is P , and hence exactly d edges meet at each vertex. At each principal vertex of B , d edges also meet. Hence, B contains every edge of P through z_0 or z_1 . Let B_0 be the subgraph of B consisting of these edges.

In B_0 two paths join z_0 and z_1 , namely those through (x_0, y_1) and (x_1, y_0) . At most one of these paths can correspond to an edge of C_{d+1} , and so at least one contains another principal vertex. So, let us suppose that (x_0, y_1) is principal. As before, B must contain all the edges of P through (x_0, y_1) ; let B_1 consist of these edges and those of B_0 . Now, z_0 and (x_0, y_1) are joined by an edge of B_1 , which must correspond to an edge of C_{d+1} . Thus every other path in B_1 joining z_0 and (x_0, y_1) must contain a principal vertex, and so the vertices (x_0, y_j) ($j = 2, \dots, q$) are all principal.

In an exactly similar way, we see that all the vertices (x_i, y_1) ($i = 0, \dots, p$) are principal. We have now found $p + q + 1 = d + 1$ principal vertices; that is, we have all of them. The graph B then consists of all the edges of P meeting these principal vertices; it is easily seen to be a refinement of C_{d+1} . For, any two principal vertices (x_0, y_j) and $(x_{i'}, y_{j'})$ are joined by an edge of B , as are any two (x_i, y_1) and $(x_{i'}, y_1)$, while any pair (x_0, y_j) and (x_i, y_1) , with $j \neq 1$ and $i \neq 0$, is joined by a path of two edges of B passing through the intermediate vertex (x_i, y_j) , which is used by this pair alone.

If (x_1, y_0) is chosen as a principal vertex, we get the alternative refinement, with principal vertices (x_i, y_0) ($i = 0, \dots, p$) and (x_1, y_j) ($j = 1, \dots, q$). The two cases are distinct, but symmetrical. We also note that one case excludes the other; (x_0, y_1) and (x_1, y_0) cannot both be principal. Finally, we observe that in neither case is z_2 a principal vertex, and this establishes Theorem 2.

Theorem 2, together with Grünbaum's result, implies that for $d \geq 4$, $r(d) = 1$ or 2. Nothing further is known, although Larman and Mani suggest that $r(d) = 2$. It is tempting to generalize the fact that if $d \leq 3$, in a d -connected graph we may preassign two vertices as principal in a refinement of C_{d+1} . However, for $d \geq 4$, a d -connected graph need not contain a refinement of C_{d+1} at all. For instance, the graph of the regular icosahedron is 5-connected, yet, by Kuratowski's theorem, it contains no refinement of C_5 , and hence none of C_6 .

We observe, however, that the conjecture $r(d) = 2$ is, in some sense, a natural generalization of the fact that $G(P)$ is d -connected if P is a d -polytope. For, the d arcs in $G(P)$, which join two given vertices, form a subgraph which is a refinement of the graph H_{d+1} obtained from C_{d+1} by deleting the edges of a complete subgraph C_{d-1} . We can recover C_{d+1} from H_{d+1} by adjoining to it the edges spanned by its $d - 1$ 2-valent vertices.

We now move on to simplicial d -polytopes; the following will apply for $d \geq 3$. We first have a result from [3].

Theorem 3. *Let P be a simplicial d -polytope, and let x_0 and x_1 be two vertices of P . Then $G(P)$ contains a refinement of C_{d+1} , with x_0 and x_1 as principal vertices.*

Proof. Let F be a facet of P containing x_1 but not x_0 (see [2, 3.1.8]), and let $\text{vert } F = \{x_1, \dots, x_d\}$. Then there are d arcs in $G(P)$, disjoint except at x_0 , joining x_0 to x_1, \dots, x_d , since $G(P)$ is d -connected. These arcs, together with the edges of F , then form the required refinement.

Note that the proof, which is a little different from that of [3], actually yields a rather stronger result than was asked for.

We thus have $r_s(d) \geq 2$. Examples of simplicial d -polytopes P for which $r(P) \leq 3$ are easily found.

Theorem 4. *Let $d \geq 3$. Then there is a simplicial d -polytope P , with some four vertices which cannot all be principal for any refinement of C_{d+1} in $G(P)$.*

Proof. We construct P as follows. Let T^d be a d -simplex, with $\text{vert } T^d = \{x_0, \dots, x_d\}$, and, for $i = 0, 1$, let F_i be the facet of T^d which does not contain x_i . Let P be obtained from T^d by erecting over each F_i a shallow pyramid with apex y_i .

Suppose that C is a refinement of C_{d+1} in $G(P)$, with y_0 and y_1 both principal. Since C_{d+1} is d -valent at each vertex, and y_0 and y_1 are d -valent vertices of $G(P)$, every edge of $G(P)$ which meets y_0 or y_1 belongs to C . Hence, for $j = 2, \dots, d$, the arc which joins y_0 and y_1 , and passes through x_j , lies in C . At most one of these arcs corresponds to an edge of C_{d+1} in the refinement, and so $d - 2$ of these arcs each contain another principal vertex, which must come from x_2, \dots, x_d . Together with y_0 and y_1 , these account for d principal vertices. Since there are only $d + 1$ principal vertices in all, we see that x_0 and x_1 cannot both be principal. Thus x_0, x_1, y_0 and y_1 cannot all be principal in any refinement of C_{d+1} in $G(P)$, which proves the theorem.

Combining Larman and Mani's result with Theorem 4 shows that for $d \geq 4$, $r_s(d) = 2$ or 3 .

References

- [1] B. Grünbaum, On the facial structure of convex polytopes, Bull. Amer. Math. Soc. 71 (1965) 559-560.
- [2] B. Grünbaum. Convex Polytopes. (Wiley-Interscience, London-New York-Sydney, 1967).
- [3] D.G. Larman and P. Mani, On the existence of certain configurations within graphs and the 1-skeletons of polytopes, Proc. London Math. Soc. 20 (3) (1970) 144-160.